

# New Hampshire Volunteer Lake Assessment Program

## 2003 Interim Report for Harvey Lake Northwood



NHDES  
Water Division  
Watershed Management Bureau  
29 Hazen Drive  
Concord, NH 03301



# OBSERVATIONS & RECOMMENDATIONS

After reviewing data collected from **HARVEY LAKE, NORTHWOOD**, the program coordinators have made the following observations and recommendations:

We would like to thank your group for sampling your lake/pond once this summer. However, we would like to encourage your monitoring group to sample additional times each summer. Typically we recommend that monitoring groups sample three times per summer (once in June, July, and August). We understand that the number of sampling events you decide to conduct per summer will depend upon volunteer availability, your monitoring group's water monitoring goals and funding availability. However, with a limited amount of data it is difficult to determine accurate and representative water quality trends. Since weather patterns and activity in the watershed can change throughout the summer, from year to year, and even from hour to hour during a rain event, it is a good idea to sample the lake/pond at least once per month over the course of the season.

If you are having difficulty finding volunteers to help sample, or to pick-up or drop-off equipment at one of the laboratories, please give the VLAP Coordinator a call and we will try to help you work out an arrangement.

We would like to encourage your monitoring group to participate in the DES Weed Watchers program, a volunteer program dedicated to monitoring the lakes and ponds for the presence of exotic weeds. This program only involves a small amount of time during the summer months. Volunteers survey their waterbody once a month from June through September. To survey, volunteers slowly boat, or even snorkel, around the perimeter of the waterbody and any islands it may contain. Using the materials provided in the Weed Watchers Kit, volunteers look for any species that are of suspicion. After a trip or two around the waterbody, volunteers will have a good knowledge of its plant community and will immediately notice even the most subtle changes. If a suspicious plant is found, the volunteers will send a specimen to DES for identification. If the plant specimen is an exotic, a biologist will visit the site to determine the extent of the problem and to formulate a plan of

action to control the nuisance infestation. Since another lake in your town has variable milfoil, an exotic plant, it is very important to prevent or detect an infestation early in Harvey Lake.

If you would like to help protect your lake or pond from exotic plants, contact Amy Smagula, Exotic Species Program Coordinator, at 271-2248 or visit the Weed Watchers web page at [www.des.state.nh.us/wmb/exoticspecies/survey.htm](http://www.des.state.nh.us/wmb/exoticspecies/survey.htm).

### **FIGURE INTERPRETATION**

- **Figure 1 and Table 1:** The graphs in Figure 1 (Appendix A) show the historical and current year chlorophyll-a concentration in the water column. Table 1 (Appendix B) lists the maximum, minimum, and mean concentration for each sampling season that the lake/pond has been monitored through the program.

Chlorophyll-a, a pigment naturally found in plants, is an indicator of the algal abundance. Because algae are usually microscopic plants that contain chlorophyll-a, and are naturally found in lake ecosystems, the chlorophyll-a concentration measured in the water gives an estimation of the algal concentration or lake productivity. **The mean (average) summer chlorophyll-a concentration for New Hampshire's lakes and ponds is 7.02 ug/L.**

The current year data (the top graph) show that the chlorophyll-a concentration in August was **greater than** state mean.

Overall, visual inspection of the historical data trend line (the bottom graph) shows **a variable** in-lake chlorophyll-a trend, meaning that the concentration has **fluctuated** since monitoring began in 1995. In the 2004 annual report, we will be able to conduct a statistical analysis of the historical data to objectively determine if there has been a significant change in the annual mean chlorophyll-a concentration since monitoring began.

While algae are naturally present in all lakes/ponds, an excessive or increasing amount of any type is not welcomed. In freshwater lakes/ponds, phosphorus is the nutrient that algae depend upon for growth. Algal concentrations may increase with an increase in nonpoint sources of phosphorus loading from the watershed, or in-lake sources of phosphorus loading (such as phosphorus releases from the sediments). Therefore, it is extremely important for volunteer monitors to continually educate residents about how

activities within the watershed can affect phosphorus loading and lake/pond quality.

- **Figure 2 and Table 3:** The graphs in Figure 2 (Appendix A) show historical and current year data for lake/pond transparency. Table 3 (Appendix B) lists the maximum, minimum and mean transparency data for each sampling season that the lake/pond has been monitored through the program.

Volunteer monitors use the Secchi-disk, a 20 cm disk with alternating black and white quadrants, to measure water clarity (how far a person can see into the water). Transparency, a measure of water clarity, can be affected by the amount of algae and sediment from erosion, as well as the natural colors of the water. **The mean (average) summer transparency for New Hampshire's lakes and ponds is 3.7 meters.**

The current year data (the top graph) show that the in-lake transparency in August was **less than** the state mean.

Overall, visual inspection of the historical data trend line (the bottom graph) shows a **relatively stable** trend for in-lake transparency. Specifically, the transparency has ranged between **approximately 2 and 3 meters** since monitoring began in 1995. As discussed previously, in the 2004 annual report we will conduct a statistical analysis of the historical data to objectively determine if there has been a significant change in the annual mean transparency since monitoring began.

Typically, high intensity rainfall causes erosion of sediments into lakes/ponds and streams, thus decreasing clarity. Efforts should continually be made to stabilize stream banks, lake/pond shorelines, disturbed soils within the watershed, and especially dirt roads located immediately adjacent to the edge of tributaries and the lake/pond. Guides to Best Management Practices designed to reduce, and possibly even eliminate, nonpoint source pollutants, such as sediment loading, are available from DES upon request.

- **Figure 3 and Table 8:** The graphs in Figure 3 (Appendix A) show the amounts of phosphorus in the epilimnion (the upper layer) and the hypolimnion (the lower layer); the inset graphs show current year data. Table 8 (Appendix B) lists the annual maximum, minimum, and median concentration for each deep spot layer and each tributary since the lake/pond has joined the program.

Phosphorus is the limiting nutrient for plant and algae growth in New Hampshire's freshwater lakes and ponds. Too much phosphorus in a lake/pond can lead to increases in plant and algal growth over time. **The median summer total phosphorus concentration in the epilimnion (upper layer) of New Hampshire's lakes and ponds is 11 ug/L. The median summer phosphorus concentration in the hypolimnion (lower layer) is 14 ug/L.**

The current year data for the epilimnion (the top inset graph) and the hypolimnion (the bottom inset graph) show that the phosphorus concentration in August was ***slightly greater than*** the state median.

Overall, visual inspection of the historical data trend line for the epilimnion and the hypolimnion show a ***variable*** phosphorus trend, which means that the concentration has ***fluctuated, but has not continually increased or decreased***, since monitoring began.

One of the most important approaches to reducing phosphorus loading to a waterbody is to continually educate watershed residents about its sources and how excessive amounts can adversely impact the ecology and value of lakes and ponds. Phosphorus sources within a lake or pond's watershed typically include septic systems, animal waste, lawn fertilizer, road and construction erosion, and natural wetlands.

#### **TABLE INTERPRETATION**

##### **➤ Table 2: Phytoplankton**

Table 2 (Appendix B) lists the current and historical phytoplankton species observed in the lake/pond. The dominant phytoplankton species observed this year were an **unidentifiable filamentous blue-green algae, *Microcystis* (a cyanobacteria), and *Tabellaria* (a diatom).**

Phytoplankton populations undergo a natural succession during the growing season (Please refer to the "Biological Monitoring Parameters" section of this report for a more detailed explanation regarding seasonal plankton succession). Diatoms and golden-brown algae are typical in New Hampshire's less productive lakes and ponds.

An overabundance of cyanobacteria (previously referred to as blue-green algae) indicates that there may be an excessive total phosphorus concentration in the lake/pond, or that the ecology is out of balance. ***Microcystis***, a cyanobacterium, was one of the dominant species in the plankton sample. In addition, **small amounts** of the cyanobacteria ***Anabaena, Aphanizomenon, and Oscillatoria*** were also observed in the plankton sample this season. ***These species, if***

***present in large amounts, can be toxic to livestock, wildlife, pets, and humans.***

Cyanobacteria can reach nuisance levels when excessive nutrients and favorable environmental conditions occur. During September of 2003, a few lakes and ponds in the southern portion of the state experienced cyanobacteria blooms. This was likely due to nutrient loading to these waterbodies. As mentioned previously, many weeks during the Spring and Summer of 2003 were rainy, which likely resulted in a large amount of nutrient loading to surface waters.

The presence of cyanobacteria serves as a reminder of the lake's/pond's delicate balance. Watershed residents should continue to act proactively to reduce nutrient loading into the lake/pond by eliminating fertilizer use on lawns, keeping the lake/pond shoreline natural, re-vegetating cleared areas within the watershed, and properly maintaining septic systems and roads.

In addition, residents should also observe the lake/pond in September and October during the time of fall turnover (lake mixing) to document any algal blooms that may occur. Cyanobacteria (blue-green algae) have the ability to regulate their depth in the water column by producing or releasing gas from vesicles. However, occasionally lake mixing can affect their buoyancy and cause them to rise to the surface and bloom. Wind and currents tend to "pile" cyanobacteria into scums that accumulate in one section of the lake/pond. If a fall bloom occurs, please contact the VLAP Coordinator.

➤ **Table 4: pH**

Table 4 (Appendix B) presents the in-lake and tributary current year and historical pH data.

pH is measured on a logarithmic scale of 0 (acidic) to 14 (basic). pH is important to the survival and reproduction of fish and other aquatic life. A pH below 5.5 severely limits the growth and reproduction of fish. A pH between 6.5 and 7.0 is ideal for fish. The mean pH value for the epilimnion (upper layer) in New Hampshire's lakes and ponds is **6.5**, which indicates that the surface waters in state are slightly acidic. For a more detailed explanation regarding pH, please refer to the "Chemical Monitoring Parameters" section of this report.

The mean pH at the deep spot this season ranged from **6.02** in the hypolimnion to **6.60** in the epilimnion, which means that the water is ***slightly acidic***. When organic material near the lake bottom is decomposed, acidic byproducts are produced, which likely explains the lower pH (meaning higher acidity) in the hypolimnion.

Due to the presence of granite bedrock in the state and the deposition of acid rain, there is not much that can be done to effectively increase lake/pond pH.

➤ **Table 5: Acid Neutralizing Capacity**

Table 5 (Appendix B) presents the current year and historical epilimnetic ANC for each year the lake/pond has been monitored through VLAP.

Buffering capacity or ANC describes the ability of a solution to resist changes in pH by neutralizing the acidic input to the lake. The mean ANC value for New Hampshire's lakes and ponds is **6.7 mg/L**, which indicates that many lakes and ponds in the state are "highly sensitive" to acidic inputs. For a more detailed explanation, please refer to the "Chemical Monitoring Parameters" section of this report.

The Acid Neutralizing Capacity (ANC) of the epilimnion (the upper layer) continues to remain **slightly less than** the state mean. Specifically, the August ANC was **5.00 mg/L**, which indicates that the lake/pond is **critically sensitive** to acidic inputs (such as acid precipitation).

➤ **Table 6: Conductivity**

Table 6 (Appendix B) presents the current and historical conductivity values for tributaries and in-lake data. Conductivity is the numerical expression of the ability of water to carry an electric current. The mean conductivity value for New Hampshire's lakes and ponds is **62.1 uMhos/cm**. For a more detailed explanation, please refer to the "Chemical Monitoring Parameters" section of this report.

The conductivity in the lake/pond and inlets has ranged between approximately 90 and 120 uMhos/cm since monitoring began, which is **greater than** the state mean. Typically, sources of increased conductivity are due to human activity. These activities include septic systems that fail and leak leachate into the groundwater (and eventually into the tributaries and the lake/pond), agricultural runoff, and road runoff (which contains road salt during the spring snow melt). New development in the watershed can alter runoff patterns and expose new soil and bedrock areas, which could contribute to increasing conductivity. In addition, natural sources, such as iron deposits in bedrock, can influence conductivity.

We recommend that your monitoring group conduct stream surveys and storm event sampling along the inlet(s) so that we can determine what may be causing the increases.

*For a detailed explanation on how to conduct rain event and stream surveys, please refer to the 2002 VLAP Annual Report “Special Topic Article”, or contact the VLAP Coordinator.*

It is also possible that de-icing materials applied to nearby roadways during the winter months may be influencing the conductivity in the tributaries and in the lake. In New Hampshire, the most commonly used de-icing material is salt (sodium chloride). Therefore, we recommend that the inlets with elevated conductivity be sampled for chloride next season.

*For a detailed explanation on how to conduct chloride sampling, please contact the VLAP Coordinator.*

➤ **Table 8: Total Phosphorus**

Table 8 (Appendix B) presents the current year and historical total phosphorus data for in-lake and tributary stations. Phosphorus is the nutrient that limits the algae’s ability to grow and reproduce. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

**Inlet 1** and the **Outlet** were sampled for phosphorus on the August sampling event. The phosphorus concentration in the **Inlet 1** sample was **slightly elevated** (42 ug/L) and the turbidity of the sample was **very slightly elevated** (2.09 NTUs). This suggests that the stream bottom may have been disturbed while sampling or that erosion is occurring in this portion of the watershed.

When the stream bottom is disturbed, sediment, which typically contains attached phosphorus, is released into the water column. When collecting samples in the inlets, please be sure to sample where there the stream is flowing and where the stream is deep enough to collect a “clean” sample.

If you suspect that erosion is occurring in this portion of the watershed, we recommend that your monitoring group conduct stream surveys and storm event sampling along this inlet. This additional sampling may allow us to determine what is causing the elevated levels of turbidity and phosphorus.

*For a detailed explanation on how to conduct rain event and stream surveys, please refer to the 2002 VLAP Annual Report “Special Topic Article”, or contact the VLAP Coordinator.*



➤ **Table 9 and Table 10: Dissolved Oxygen and Temperature Data**

Table 9 (Appendix B) shows the dissolved oxygen/temperature profile(s) for the 2003 sampling season. Table 10 (Appendix B) shows the historical and current year dissolved oxygen concentration in the hypolimnion (lower layer). The presence of dissolved oxygen is vital to fish and amphibians in the water column and also to bottom-dwelling organisms. Please refer to the “Chemical Monitoring Parameters” section of this report for a more detailed explanation.

The dissolved oxygen concentration was **low in the hypolimnion** at the deep spot of the lake/pond. As stratified lakes/ponds age, and as the summer progresses, oxygen typically becomes **depleted** in the hypolimnion (the lower layer) by the process of decomposition. Specifically, the loss of oxygen in the hypolimnion results primarily from the process of biological breakdown of organic matter (i.e.; biological organisms use oxygen to break down organic matter), both in the water column and particularly at the bottom of the lake/pond where the water meets the sediment. When oxygen levels are depleted to less than 1 mg/L in the hypolimnion (**as it was this season and in many past seasons**), the phosphorus that is normally bound up in the sediment may be re-released into the water column.

The **low** oxygen level in the hypolimnion is a sign of the lake’s/pond’s **aging** and **declining** health. This year the DES biologist conducted the temperature/dissolved oxygen profile in **August**. We recommend that the annual biologist visit for the 2004 sampling season be scheduled during **June** so that we can determine if oxygen is depleted in the hypolimnion **earlier** in the sampling season.

➤ **Table 11: Turbidity**

Table 11 (Appendix B) lists the current year and historical data for in-lake and tributary turbidity. Turbidity in the water is caused by suspended matter, such as clay, silt, and algae. Water clarity is strongly influenced by turbidity. Please refer to the “Other Monitoring Parameters” section of this report for a more detailed explanation.

## **DATA QUALITY ASSURANCE AND CONTROL**

### **Annual Assessment Audit:**

Since this was the first time your group sampled the lake, the biologist trained your group how to collect samples during the annual biologist visit. Therefore, the biologist did not conduct a “Sampling Procedures Assessment Audit” for your monitoring group. In future sampling seasons, the biologist will observe the performance of your monitoring group while sampling and will fill out an assessment audit sheet to

document the ability of the volunteer monitors to follow the proper field sampling procedures (as outlined in the V LAP Monitor's Field Manual). This assessment is used to identify any aspects of sample collection in which volunteer monitors are not following the proper procedures, and also provides an opportunity for the biologist to retrain the volunteer monitors as necessary. This will ultimately ensure that the samples that the volunteer monitors collect are truly representative of actual lake and tributary conditions.

### **USEFUL RESOURCES**

*Acid Deposition Impacting New Hampshire's Ecosystems*, ARD-32, NHDES Fact Sheet, (603) 271-3505, or [www.des.state.nh.us/factsheets/ard/ard-32.htm](http://www.des.state.nh.us/factsheets/ard/ard-32.htm).

*Aquarium Plants and Animals: Don't leave them stranded*. Informational pamphlet sponsored by NH Fish and Game, Aquaculture Education and Research Center, and NHDES (603) 271-3505.

*Best Management Practices to Control Nonpoint Source Pollution: A Guide for Citizens and Town Officials*, NHDES-WD 97-8, NHDES Booklet, (603) 271-3503.

*A Boater's Guide to Cleaner Water*, NHDES pamphlet, (603) 271-3503.

*Camp Road Maintenance Manual: A Guide for Landowners*. Kennebec Soil and Water Conservation District, 1992, (207) 287-3901.

*Comprehensive Shoreland Protection Act*, RSA 483-B, WD-SP-5, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/sp/sp-5.htm](http://www.des.state.nh.us/factsheets/sp/sp-5.htm).

*Cyanobacteria in New Hampshire Waters Potential Dangers of Blue-Green Algae Blooms*, NHDES Fact Sheet, (603) 271-3505, or [www.des.state.nh.us/factsheets/wmb/wmb-10.htm](http://www.des.state.nh.us/factsheets/wmb/wmb-10.htm).

*Erosion Control for Construction in the Protected Shoreland Buffer Zone*, WD-SP-1, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/sp/sp-1.htm](http://www.des.state.nh.us/factsheets/sp/sp-1.htm)

*Impacts of Development Upon Stormwater Runoff*, WD-WQE-7, NHDES Fact Sheet, (603) 271-3503, or [www.des.state.nh.us/factsheets/wqe/wqe-7.htm](http://www.des.state.nh.us/factsheets/wqe/wqe-7.htm)

*Iron Bacteria in Surface Water*, WD-BB-18, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/bb/bb-18.htm](http://www.des.state.nh.us/factsheets/bb/bb-18.htm)

*Is it Safe to Eat the Fish We Catch? Mercury and Other Pollutants in Fish*, NH Department of Health and Human Services pamphlet, 1-800-852-3345, ext. 4664.

*Lake Protection Tips: Some Do's and Don'ts for Maintaining Healthy Lakes*, WD-BB-9, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/bb/bb-9.htm](http://www.des.state.nh.us/factsheets/bb/bb-9.htm).

*Management of Canada Geese in Suburban Areas: A Guide to the Basics*, Draft Report, NJ Department of Environmental Protection Division of Watershed Management, March 2001, [www.state.nj.us/dep/watershedmgt/DOCS/BMP\\_DOCS/Goosedraft.pdf](http://www.state.nj.us/dep/watershedmgt/DOCS/BMP_DOCS/Goosedraft.pdf).

*Proper Lawn Care In the Protected Shoreland, The Comprehensive Shoreland Protection Act*, WD-SP-2, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/sp/sp-2.htm](http://www.des.state.nh.us/factsheets/sp/sp-2.htm).

*Road Salt and Water Quality*, WD-WMB-4, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/wmb/wmb-4.htm](http://www.des.state.nh.us/factsheets/wmb/wmb-4.htm).

*Sand Dumping - Beach Construction*, WD-BB-15, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/bb/bb-15.htm](http://www.des.state.nh.us/factsheets/bb/bb-15.htm).

*Swimmers Itch*, WD-BB-2, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/bb/bb-2.htm](http://www.des.state.nh.us/factsheets/bb/bb-2.htm).

*Through the Looking Glass: A Field Guide to Aquatic Plants*. North American Lake Management Society, 1988, (608) 233-2836 or [www.nalms.org](http://www.nalms.org).



*Weed Watchers: An Association to Halt the Spread of Exotic Aquatic Plants*, WD-BB-4, NHDES Fact Sheet, (603) 271-3503 or [www.des.state.nh.us/factsheets/bb/bb-4.htm](http://www.des.state.nh.us/factsheets/bb/bb-4.htm).

# APPENDIX A

## GRAPHS



# Harvey Lake, Northwood

**Figure 1.** Monthly and Historical Chlorophyll-a Results



## Harvey Lake, Northwood

**Figure 2.** Monthly and Historical Transparency Results



# Harvey Lake, Northwood

**Figure 3.** Monthly and Historical Total Phosphorus Data.

